

ECAT Chemistry Chapter 5 Atomic Structure Online Test

Sr	Questions	Answers Choice
1	Which of the following is not a property of cathode rays	 A. They can produce x-rays when they strike a heavy metal anode B. They can cause reduction reaction C. They produce fluorescence in rare earth and minerals D. They comprise neutral particles
2	Which of the following is not a sub-atomic particle	A. Electron B. Proton C. Neutron D. Deuteron
3	Which of the following is not charged particle	A. Proton B. Electron C. Neutron D. Hydrogen nucleus
4	The radius of first orbit of hydrogen atom	A. 0.329 A° B. 0.429 A° C. 0.529 A° D. 0.229 A°
5	Splitting of spectral lines when atoms are subjected to strong electron field is called	A. Zeeman effectB. Stark effectC. Photoelectric effectD. Compton effect
6	E = hv is the	 A. Spectral equation B. Plank's equation C. de Broglie's equation D. None of these
7	The nature of the positive rays depend on	A. The nature of the electrode B. The nature of the discharge tube C. The nature of the residual gas D. All of the above
8	The wave number of the light emitted by a certain source is 2 x 10^{6} m ⁻¹ . The wavelength of this light will be	A. 500 nm B. 500 m C. 200 nm D. 5 x 10 ⁷ m
9	Pauli's principle is applicable to	A. Degenerate orbits B. Two electrons in the same orbital C. One electron D. None
10	The charge of an electron is determined by	A. J.J. Thomson B. Crooks C. Perrin D. R.A.Millikan
11	Quantum number values for 2p orbitals are	A. n = 2, I = 1 B. n = 1, I = 2 C. n = 1, I = 0 D. n = 2, I = 0
12	In the ground state of an atom, the electron is present	A. In the nucleus B. In the second shell C. Nearest to the nucleus D. Farthest from the nucleus
13	When 3p orbital is complete, the entering electron goes into	A. 4s B. 3d C. 4p D. 4f
14	Orbitals having same energy are called	A. Hybrid orbitals B. Valence orbitals C. Degenerate orbitals D. D-orbitals
15	The wave length of electron as wave is 0.5 nm. What is the wave length in meter	A. 5 x 10 ⁻⁹ B. 5 x 10 ⁻¹² C. 5 x 10 ⁻⁶

		D. 5 x 10 ⁻¹⁰
16	The principle quantum number describes	A. The distance form the nucleusB. The shape of the orbitalC. The orientation of the orbitalD. The spin of the electron
17	The rules which describe the distribution of electron in atomic energy levels are Auf-ban principle, Pauli's exclusion principle. Hunds rule. The pauli exclusion principle refers to the	 A. Orientation of orbital in space B. Fact that two electrons in the same orbital should have opposite spins C. Energy of the orbital D. Spin of the electron
		A. s
18	Question Image	C.d D.f
19	The value of charge on electron is	A. 1.602 x 10 ⁻¹⁹ coulombs B. 1.602 x 10 ⁻¹⁸ coulombs C. 1.602 x 10 ⁻¹⁷ coulombs D. 1.602 x 10 ⁻¹⁶ coulombs
20	Cathode rays drive a small paddle wheel placed in their path. This observation shows that	 A. Cathode rays travel in straight lines B. Cathode rays are negatively charged C. Cathode rays produce x-rays D. Cathode rays are material particles having momentum
21	No cathode rays are produced in the discharged tube when gas is under ordinary pressure even if voltage of 5000 to 10000 is applies. This reason is	A. Voltage is low B. Discharge tube is not coloured C. Gas does not conduct current under ordinary pressure D. Temperature low
22	e/m of cathode rays is same but for positive rays e/m changes by changing gas in the discharge tube because	 A. Cathode rays are small sized particles B. Cathode rays have same charge C. Nature of cathode rays same for all gasses, but masses of nuclei are different for different gases D. Temperature of cathode rays higher
23	When cathode rays strike the anode metal X-rays are emitted and not the positive rays because	 A. Cathode rays are material particles B. Cathode rays knock out electrons from anode, which emit X-rays when outer electron take their place C. Cathode rays are absorbed by the nucleus D. Cathode rays become heated
24	Positive particle in discharged tube is produced by ionization of gas molecules, which is caused by	A. Gas molecules collide with anode B. Gas molecules are at high temperature C. Gas molecules produce X-rays D. Cathode rays remove electrons from gas molecules
25	When the 6d orbital is completed the entering electron goes into	A. 7f B. 7 s C. 7 p D. 7 d
26	Question Image	 A. Neutrons are attracted by nucleus B. Neutrons carry out nuclear reactions C. Neutrons carry no charge D. Neutrons are electromagnetic radiations
27	In Millikan method for determination of charge on electron the air in the chamber is ionized by	A. Protons B. Electric field C. X-rays D. a - particles
28	In Millikan method the oil droplet falls under the force of gravity but it moves upward due to	A. Electric field B. Magnetic field C. Incident light D. X-rays
29	Smallest charge of electricity that has been measured so far is	A. Charge on a-rays B. Charge on electron (1.602 x 10 ^{- 19} C) C. Charge on x-rays D. Charge on gamma rays
30	Rutherford's planetary like picture of the atomic modal was defective because	 A. It did not describe the quantity of positive charge B. It did not explain the repulsion of protons within the nucleus C. No empty space between nucleus and the electrons D. Moving electron should radiate energy
		A. r ₂ - r ₁ > r ₃ - r ₂ >

31	The order of distance between the various Bohr orbits is	<pre>rsub>3- rsub>2> B. rsub>1> rsub>3> B. rsub>1> rsub>3> rsub>2- rsub>3> rsub>4-rsub>3> C. rsub>2-rsub>1= rsub>3- rsub>2= rsub>1- rsub>3- rsub>2- rsub>2- rsub>4- rsub>3> D. rsub>2- rsub>2> rsub>3- rsub>3<!--</th--></pre>
32	The radius of first orbit of H-atom is	A. 4.75 A° B. 3.84 A° C. 8.4 A° D. 0.529 A°
33	Energy of electron in first orbit of H atom is	A45.32 KJ/mole B82.08 KJ/mole C52.53 KJ/mole D1313.31 KJ/mole
34	Energy of electron in the infinite Bohr orbit of H-atom is	A. 0 KJ/mole B. 1 KJ/mole C1 KJ/mole D1313.32 KJ/mole
35	In Bohr model of hydrogen atom the distance between adjacent orbits increases away from the nucleus, the energy difference between the orbits	A. Increases B. Decreases C. Reaming same D. Orbits coincide
36	Energy of electron in an orbit according to Bohr theory is negative due to	 A. Repulsion of electrons in the same orb B. At infinity energy is zero ad a traction towards nucleus decreases energy C. Electron has negative charge D. Product of positive nuclear charge and negative charge is negative
37	In the atomic emission spectrum the lines which appear bright, appear dark in absorption spectrum because	 A. The radiations emitted in emission spectrum are absorbed in absorption spectrum B. Atomic emission spectrum is continuous C. Atomic absorption spectrum is continuous D. Distance between the lines increases
38	When electron jumps from n_2 = 2,3,4,5, orbit to n_1 = 1 orbit in the hydrogen atom, the radiations emitted give the spectral lines	A. Lyman series B. Blamer series C. Paschen series D. Brackett series
39	The limiting line of Blamer series in hydrogen spectrum lies in	A. Visible regions B. Ultraviolet region C. Infrared region D. x-rays region
40	Splitting of spectral lines of the hydrogen atom under the influence or magnetic field is called	A. Stark effect B. Zeeman effect C. Compton effect D. Photoelectric effect
41	In the ground state of an atom the electron is present	A. In the nucleusB. In the second shellC. Nearest to the nucleusD. Farthest from the nucleus
42	Light emitted from a source has its wave length 500nm, then its wave number will be	A. 2 x 10 ⁶ m ⁻¹ B. 2 x 10 ⁷ m ⁻¹ C. 5 x 10 ⁸ m ⁻¹ D. 5 x 10 ⁹ m ⁻¹
43	The wave number of the line emitted is 109.678 x $10^{5}m^{-1}$ in the Lyman series when electron transition occurs	
44	The divisibility of atom was shown by	A. Stoney B. J.J. Thomson C. Millikan D. Rutherford
45	Question Image	A. Plank's equations B. de Broglie's equations C. Heisenburg's equation D. None
46	The uncertainty principle was stated only	A. De Brogilie B. Heinsenberg C. Einstein D. Schrodinger
		A. 6.625 x 10 ⁻³⁴

47	The value of Plank's constant 'h' is	B. 6.625 x 10 ⁻³⁴ J sec C. 6.625 x 10 ⁻³⁴ KJ D. 6.625 x 10 ⁻³⁴ KCal
48	Schrodinger wave equation describes electron completely because	 A. It describes a set of four quantum number B. It describes the particle nature of electron C. It measures wavelength of electron D. It describes electron moving in specific orbit
49	The size of electronic shell is described by	A. Azimuthal Q. no B. Magnetic Q.No C. Spin Q. No D. Principle Q. No
50	The total values of magnetic quantum number of subshell are five, the subshell is	A. S-subshell B. P-subshell C. D-subshell D. F-subshell
51	For a 3P subshell the set of principle and azimuthal quantum number is	A. n = 1 , l = 2 B. n = 3, l = 0 C. n = 3, l = 1 D. n = 1, l = 3
52	The orbitals having n + I = 5 are	A. 2p, 3d,3s B. 3p, 3d, 5s C. 3s, 4p, 4d D. 5s, 4p, 3d
53	Their e/m, ratio resembles with that of electrons	A. Alpha rays B. Beta rays C. Gamma rays D. X-rays
54	An orbital can accommodate maximum two electrons with opposite spins according to	A. Heisenberg's principle B. Aufbau principle C. Hund's srule D. Pauli exclusion principle
55	The degenerate orbitals p-sub shell are	A. 2 B. 3 C. 5 D. 7
56	Electrons arranged in orbitals according to the increasing order of their n + I values, this rule is named as	A. Hund's ruleB. Heisenberg's principleC. Paulit exclusion principleD. Auf bau principle
57	The arrangement of subshells in the ascending order of their energy on complete filing of 4f subshell the entering electrons goes to	A. 5s B. 5p C. 5d D. 5f
58	When 6s orbital is complete then next electron goes to	A. 6p B. 6d C. 5d D. 4f
59	Four d-orbitals contain four lobes while fifth contains only two lobes the orbital is	A. dxy B. dxz C. dz ² D. dx ² - y ²
60	The energy of ionization of an atom is the energy difference between orbital	
61	Spectrum of white light is continuous becuase	 A. Colors separated by dark spaces B. There are no boundary lines between the colours C. The radiations are in infrared region D. The radiatins fall in ultraviolet region
62	The radiations with wavelength shorter than violet light are called	A. Ultraviolet B. Infrared C. Microwave D. Radio frequency
63	Which have better penetrating power	A. Alpha rays B. Beta rays C. Gamma rays D. X-rays
64	The order of frequency of the following radiations unltraviolet, visible, infrared and microwave is	 A. Microwave > infrared > visible > ultraviolet B. Visible > ultraviolet > microwave > infrared C. Ultraviolet > visible > infrared > microwave D. Infrared > microwave > ultraviolet >

		visible
65	If the value of azimuthal quantum number is 3, then values of m the magnetic quantum no. will be	A. 0, 1, 2, 3 B. +3, +2, +1, -1, -2, -3 C. 0, -1, -2, -3 D3, 0, +3
66	An electron with $n = 3$, $I = 2$ will be in the sub-shell	A. 3p B. 3d C. 3f D. 3s
67	Electrons in degenerate orbitals are placed in separate orbitals with same spin according to	A. Hund's rule B. Pauli exclusion principle C. Aufbau principle D. Mosley's law
68	Which of the atoms has $1s^2$, $2s^2$, $2p_x^2 2p_y^1 2p_z^1 2p_z^1$	A. Nitrogen B. Carbon C. Fluorine D. Oxygen
69	The value of R (Rydberg's constant) is m-1	A. 1.0974 x 107 B. 1.0842 x 107 C. 1.082 x 10-7 D. Both a and b
70	Cathode rays emitted from cathode are	A. Canal rays B. Protons C. Electrons D. Positrons
71	Charge to mass ratio (e/m) of the electron is determined by	A. R. A. Millikan B. J. J. Thompson C. G. J, Stoney D. None of these
72	Neutrons was discovered by	A. Mosely B. Milliken C. Chadwick D. Ruherford
73	n + I value for 4f will	A. 2 B. 5 C. 7 D. 9
74	The range of visible spectrum is	A. 300 - 600 nm B. 600 - 900 nm C. 400 - 750 nm D. 100 - 300 nm
75	Balmer's series is in region	A. Visible B. U V C. I. R. D. None
76	The quantum number which describes the shape of the orbital is	 A. Principle quantum number B. Spin quantum number C. Azimathal quantum number D. Magnetic quantum number
77	The quantum number which describe the orientation of the orbitals is	A. Spin quantum numberB. Principle quantum numberC. Azimathal quantum numberD. Magnetic quantum number
78	Photons of yellow colour are energetic than violet colour	A. More B. Less C. Equal D. None
79	When the electron jumps form third, fourth, fifth orbits to the second orbit, the transitions are known as	A. Paschen B. Pfund C. Balmer D. Brackett
80	When the electron jumps form second third, fourth orbit to the fist orbit, the transitions are known as	A. Balmer series B. Lyman series C. Pfund series D. Brackett series
81	With increasing principle quantum number, the energy difference between adjacent energy levels in H atom	A. Decreases B. Increases C. Remains constant D. Decreases for low value of Z and increase for higher value of Z

82	The credit of discovering neutron goes to	A. Rutherford B. Langmuir C. Chadwick D. Austen
83	The mass of the neutron is of the order of	A. 10 ⁻²³ kg B. 10 ⁻²⁴ kg C. 10 ⁻²⁶ kg D. 10 ⁻²⁷ kg
84	The ratio of the ionization energy of H and Be^{3+} is	A. 1 : 1 B. 1 : 3 C. 1 : 9 D. 1 : 16
85	The maximum number of electrons in a subshell for which I = 3 is	A. 14 B. 10 C. 8 D. 4
86	The number of electrons in the M shell of the element with atomic number 24 is	A. 24 B. 12 C. 13 D. 8
87	The symbol of the element whose atoms have the outer most electronic configuration $2s^22p^3$ is	A. N B. Li C. P D. Na
88	When electrons revolve in stationary orbits	A. There is no change in energy levelB. They vecome stationaryC. They are gaining kinetic energyD. There is increase in energy
89	Which quantum number is sufficient to describe the electron is hydrogen atom?	A. I B. n C. m D. s
90	The valence orbital configuration of an element with atomic number 23 is	A. 3d ⁵ B. 3d ³ , 4s ² C. 3d ³ , 4s ¹ , 4p ¹
		3d ² ,4s ² ,4p ¹
91	The number of neutrons in the element ⁹ 4Be is	D. 3d ² ,4s ² ,4p ¹ A. 4 B. 5 C. 9 D. 13
91 92	The number of neutrons in the element ⁹ ₄ Be is Sodium chloride imparts a yellow colour to the Bunsen flame. This can be interpreted due to the	 3d²,4s²,4p¹ A. 4 B. 5 C. 9 D. 13 A. low ionization energy of sodium B. sublimation of metallic sodium to give yellow vapour C. emission of excess energy absorbed as a radiation in the visible region as a radiation in the visible region D. photosenitivity
91 92 93	The number of neutrons in the element ⁹ ₄ Be is Sodium chloride imparts a yellow colour to the Bunsen flame. This can be interpreted due to the Subsidiary quantum number specifies	A. 4 B. 5 C. 9 D. 13 A. low ionization energy of sodium B. sublimation of metallic sodium to give yellow vapour C. emission of excess energy absorbed as a radiation in the visible region as a radiation in the visible region D. photosenitivity A. size of orbital B. shape of orbital C. orientations of orbitals D. Nuclear stability
91 92 93 94	The number of neutrons in the element ⁹ 4Be is Sodium chloride imparts a yellow colour to the Bunsen flame. This can be interpreted due to the Subsidiary quantum number specifies The spectrum of helium is expected to be similar to that of	A. 4 B. 5 C. 9 D. 13 A. low ionization energy of sodium B. sublimation of metallic sodium to give yellow vapour C. emission of excess energy absorbed as a radiation in the visible region as a radiation in the visible region D. photosenitivity A. size of orbital B. shape of orbital C. orientations of orbitals D. Nuclear stability A. H B. Li ⁺ C. Na D. He ⁺
91 92 93 94 95	The number of neutrons in the element ⁹ 4Be is Sodium chloride imparts a yellow colour to the Bunsen flame. This can be interpreted due to the Subsidiary quantum number specifies The spectrum of helium is expected to be similar to that of The number of spherical nodes in 3p orbitals are	A. 4 B. 5 C. 9 D. 13 A. low ionization energy of sodium B. sublimation of metallic sodium to give yellow vapour C. emission of excess energy absorbed as a radiation in the visible region as a radiation in the visible region D. photosenitivity A. size of orbital B. shape of orbital C. orientations of orbitals D. Nuclear stability A. H B. Li ⁺ C. Na D. He ⁺ C. Na D. He ⁺ A. One B. Three C. Non D. Two
91 92 93 94 95 96	The number of neutrons in the element ⁹ 4Be is Sodium chloride imparts a yellow colour to the Bunsen flame. This can be interpreted due to the Subsidiary quantum number specifies The spectrum of helium is expected to be similar to that of The number of spherical nodes in 3p orbitals are Number of neutrons in heavy hydrogen atom is	A. 4 B. 5 C. 9 D. 13 A. low ionization energy of sodium B. sublimation of metallic sodium to give yellow vapour C. emission of excess energy absorbed as a radiation in the visible region as a radiation in the visible region D. photosenitivity A. size of orbital B. shape of orbital C. orientations of orbitals D. Nuclear stability A. H B. Li ⁺ C. Na D. He ⁺ C. Na D. He ⁺ A. One B. Three C. Non D. Two
91 92 93 94 95 96 97	The number of neutrons in the element ⁹ 4Be is Sodium chloride imparts a yellow colour to the Bunsen flame. This can be interpreted due to the Subsidiary quantum number specifies The spectrum of helium is expected to be similar to that of The number of spherical nodes in 3p orbitals are Number of neutrons in heavy hydrogen atom is The electronic configuration of an atom/ion can be defined by the following	A. 4 B. 5 C. 9 D. 13 A. low ionization energy of sodium B. sublimation of metallic sodium to give yellow vapour C. emission of excess energy absorbed as a radiation in the visible region as a radiation in the visible region D. photosenitivity A. size of orbital B. shape of orbital C. orientations of orbitals D. Nuclear stability A. H B. Li ⁺ C. Na D. He ⁺ A. One B. Three C. Non D. Two A. 0 B. 1 C. 2 D. 3 A. Aufbau principle B. Pauli's exclusion principle C. Hund's Rule D. All the above

		D. 1
99	If the value of principal quantum number is 3. the total possible values for magnetic quantum number will be	A. 1 B. 4 C. 9 D. 12
100	The electron in an atom	 A. moves randomly around the nucleus B. has fixed space around the nucleus C. is stationary in various energy levels D. moves around its nucleus in definite energy levels
101	The radius of second Bohr's orbit is	A. 0.053 nm B. 0.053/4 nm C. 0.053 x 4 nm D. 0.053 x 20 nm
102	For which of the following sets of quantum numbers and electron will have the highest energy?	A. 3,2,1,1/2 B. 4,2,-1,1/2 C. 4,1,0,-1/2 D. 5,0,0,1/2
103	Azimuthal quantum number of last electron of ₁₁ Na is	A. 1 B. 2 C. 3 D. 0
104	Rutherford's experiment led to the discovery of	A. Nucleus B. Electron C. Proton D. alpha particle
105	The total number of orbitals in a shell with principal quantum number 'n' is	A. 2n B. 2n ² C. n ² D. n + 1
106	Which of the following orbitals have a dumb bell shape?	A. s B. p C. d D. f
107	The quantum number which determines the shape of the orbital is	A. principal B. azimuthal C. magnetic D. spin
108	If the radius of first Bohr orbit be \mathbf{a}_0 , then the radius of third Bohr orbit would be	A. 3 x a _o B. 6 x a _o C. 9 x a _o D. 1/2 x a _o
109	An electron has principal quantum number 3. The number of its 1 subshell and 2 orbitals would be respectively	A. 3 and 5 B. 3 and 7 C. 3 and 9 D. 2 and 5
110	The four quantum numbers of the valency electron of potassium are	A. 4,1,1,1/2 B. 4,0,0,1/2 C. 4,1,0,1/2 D. 4,4,0,1/2
111	Heaviest particle is	A. Meson B. Neutron C. Proton D. Electron
112	Which of the following has more unpaired d-electrons?	A. Zn ⁺ B. Fe ²⁺ C. Ni ³⁺ D. Cu ⁺
113	The de-Brogile wavelength of a particle with mass 1g and velocity 100 m/s is	A. 6.63 x 10 ⁻³³ m B. 6.63 x 10 ⁻³⁴ m C. 6.63 x 10 ⁻³⁵ m D. 6.65 x 10 ⁻³⁵ m
114	1 erg of energy corresponds to	A. 6.02 x 10 ²³ J/mol B. 6.02 x 10 ¹⁶ J/mol C. 1 erg/mol D. 10 ⁻⁷ J/mol
115	A 4f orbital has	A. one node B. two node C. three node D. four nodes

A. n

116	The total number of orbitals possible for the quantum number n is	B. n ² C. 2n D. 2n ²
117	In which of the following pairs, the numbers of electrons in the outermost shell are different?	A. As,Sb B. Ge,Sn C. In,pt D. Se,Te
118	The third line of the Balmer series, in the emission spectrum of the hydrogen atom, is due to the transition from the	 A. Fourth Bohr orbit to the first Bohr orbit B. Fifth Bohr orbit to the second Bohr orbit C. Sixth Bohr orbit to the third Bohr orbit D. Seventh Bohr orbit to the third Bohr orbit
119	Rutherford's atomic model suggests the existence of	A. Atom B. Nucleus C. alpha particle D. Measons
120	In the ground state, an element has 13 electrons in its M shell. The element is	A. Copper B. Chromium C. Nickel
121	Which is not true with respect to cathode rays?	D. Iron A. A stream of electrons B. Charged particles C. Move with speed as that of light D. Can be deflected by magnetic fields
122	Which of the following element's outermost orbits last electron has magnetic quantum number m=0?	A. Na B. O C. Cl D. N
123	The spectrum of He is expected to be similar to that of	A. H B. Na C. He ⁺ D. Li ⁺
124	Which of the following is not isoelectronic?	A. Na ⁺ B. Mg ²⁺ C. O ²⁻ D. Cl ⁻
125	The energy of the first electron is helium will be	A13.6 eV B54.4 eV C5.44 eV D. zero
126	The configuration 1s ² 2s ² 2p ⁵ 3s ¹ shows	 A. Excited state of O₂ B. Excited state of neon C. Excited state of fluorine D. Ground state of fluorine atom
127	The number of de-electrons retained in Fe^{2+} (At.No. of Fe = 26) ions is	A. 3 B. 4 C. 5 D. 6
128	The orbital in Rutherford's model is	A. Spiral B. Circular C. Both D. None
129	The atomic number of an element is 35 what is the total number of electrons present in all the p-orbitals of the ground state atom of that element?	A. 6 B. 11 C. 17 D. 23
130	For principle quantum number n=4, the total number of orbitals having $I = 3$ is	A. 3 B. 7 C. 5 D. 9
131	The correct set of quantum numbers (n,I and m) respectively of the unpaired electron of chlorine atom is	A. 2,1,0 B. 2,1,1 C. 3,1,1 D. 3,2,1
132	Consider the ground state of Cr atom (Z=24). The numbers of electrons with the azimuthal quantum numbers I = 1 and I = 2 are respectively	A. 12 and 4 B. 16 and 5 C. 16 and 4 D. 12 and 5
133	Effective magnetic moment of Sc ³⁺ ion is	A. 1.73 B. 0 C. 5.92 D. 2.83

134	What is the packet of energy called?	A. Electron B. Photon C. Positron D. Proton
135	Find the magnetic moment of a divalent ion in aqueous solution if its atomic number is 25	A. 3.0 BM B. 4.9 BM C. 5.9 BM D. 6.9 BM
136	When potassium metal is exposed to violet light	 A. Ejection of electrons takes place B. Ejection of some potassium atoms occurs C. There is no effect D. The absorption of electrons takes place
137	The nature of positive ray depend on:	A. The nature of electrode.B. The nature of discharge tube.C. The nature of residual gas.D. All of above.
138	The velocity of photon is:	A. Independent of its wavelength.B. Depends on its wavelength.C. Equal to square of its amplitudeD. Depends on its source.
139	The wave number of light emitted by a certain source is 2 x $10^5 \text{m}.$ The wavelength of this light will be:	A. 500 NM. B. 500 M. C. 200 NM. D. 5 x 10 ⁷ m
140	Rutherford's model of atom failed because:	 A. The atom did o have a nucleus and electrons B. It did not account fro the attraction b/w protons and neutrons C. It did no account for the stability of the atom. D. There is actually no space b/w the nucleus ad the electrons.
141	Bohr model of atom is contradicted by:	 A. Planck's quantum theory B. Pauli's exclusion theory C. Heisenberg's uncertainty principle. D. All of above.
142	Splitting of spectral lines when atoms are subjected to strong electric field is called:	 A. Zeeman effect. B. Stark effect C. Photoelectric effect. D. Compton effect.
143	In the ground state of an atom, the electron is present:	A. In the nucleus.B. In the second shell.C. Nearest to the nucleus.D. farthest from the nucleus.
144	Quantum number value for 2p orbitals are:	A. n=2 ,I=1 B. n=1 ,I=2 C. n= ,I=0 D. n=2 ,I=0
145	Orbital having same energy is called:	A. Hybrid orbital. B. Valence orbital. C. Degenerate orbital. D. D-orbital.
146	When 6d orbital is complete, the entering electron goes into:	A. 7f. B. 7s. C. 7p. D. 7d.
147	Neutron was discovered by:	A. Chadwick. B. Bohr. C. Rutherford. D. Plank.
148	The e.m value for positive rays maximum for:	A. Oxygen. B. Nitrogen. C. Helium. D. Hydrogen.
149	The nature of positive rays depend on:	A. Nature of discharge tube.B. Nature of resident gas.C. Nature of electrode.D. All of above.
150	Which of the following was discovered first:	A. Charge to mass ratio of electrons.B. Mass of electrons.C. Charge of electrons.D. All of above at same time.
		A. Proton.

151	Which of the following particles has longest wavelength, if they have same speed:	B. Neutron. C. Electron. D. Positron.
152	Maximum potential energy that an electron can have within the atom is:	A. Equal to zero.B. Less than zero.C. Greater than zero.D. Infinite
153	Alpha rays consist of:	A. Neutrons. B. Helium nucleus. C. Protons. D. Hydrogen nucleus.
154	Charge to mass ratio of electron was discovered by:	A. Millika. B. Rutherford. C. J.J. Thomson. D. Chadwick.
155	When an electric current is passed through discharge tube at low pressure, cathode rays are emitted from cathode these rays consist of:	A. Alpha rays.B. Negative particles.C. Electromagnetic rays.D. Positive particles.
156	Charge of an electron is:	 A. <span style="font-size:
10.5pt; line-height: 107%; font-family: Arial, sansserif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-repeat: initial; background-clip: initial; background-repeat: initial; background-clip: initial; ">1.6 x 10⁻¹⁹ C<o:p></o:p> B. 9.1 x <span font-size:<br="" style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-position: initial; background-size: initial; background-position: initial; background-size: initial; background-repeat: initial; background-clip: initial; background-origin: initial; background-clip: initial; background-origin: initial; background-clip: initial; background-position: initial; background-image: initial; background-repeat: initial; background-attachment: initial; background-position: initial; background-attachment: initial; background-repeat: initial; background-attachment: initial; background-repeat: initial; background-attachment: initial; background-repeat: initial; background-clip: initial; background-repeat: initial; background-clip: initial; background-repeat: initial; background-clip: initial; background-image: initial; background-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-image: initial; background-position: initial; background-size: initial; background-position: initial; background-size: initial; background-image: initial; background-position: initial; background-size: initial; background-size: initial; background-size: initial; background-size: initial; background-size: initial; background-size: ini</td></tr><tr><td>157</td><td>Mass of simple electron is:</td><td> A. 10.5pt; line-height: 107%; font-family: Arial, sans- serif; background-image: initial; background- position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">9.1 x 10⁻³¹ kg<o:p> </o:p> B. <span style="font-size:
10.5pt; line-height: 107%; font-family: Arial, sans-
serif; background-image: initial; background-
position: initial; background-size: initial;
background-repeat: initial; background-
position: initial; background-size: initial;
background-repeat: initial; background-attachment:
initial; background-origin: initial; background-clip:
initial;">9.1 x 10⁻³⁰ kg<o:p> </o:p> C. <span style="font-size:
10.5pt; line-height: 107%; font-family: Arial, sans- serif; background-image: initial; background- position: initial; background-size: initial; background-repeat: initial; background- position: initial; background-size: initial; background-repeat: initial; background-clip: initial; background-reigin: initial; background-clip: initial; background-origin: initial; background-clip: initial; background-origin: initial; background-size: initial; background-position: initial; background-size: initial; background-repeat: initial; backg

		position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">9.1 x 10 ⁻³¹ kg <o:p> </o:p>
158	The charge over mass ratio of electron is:	A. <span font-size:<br="" style="font-size:
10.5pt; line-height: 107%; font-family: Arial, sans-
serif; background-image: initial; background-
position: initial; background-size: initial;
background-repeat: initial; background-attachment:
initial; background-origin: initial; background-clip:
initial; background-origin: initial; background-clip:
initial; background-size: initial; background-size:
10.5pt; line-height: 107%; font-family: Arial, sans-
serif; background-image: initial; background-position: initial; background-size: initial;
background-repeat: initial; background-attachment:
initial; background-origin: initial; background-attachment:
initial; background-origin: initial; background-attachment:
initial; background-origin: initial; background-attachment:
initial; '>9.1 x 10<sup>31</sup></spans C
Kg<sup>-1</sup><o:p></o:p>
C. 10.5pt; line-height: 107%; font-family: Arial, sans- serif; background-image: initial; background- position: initial; background-size: initial; background-repeat: initial; background- position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial; '>1.7588 x 10¹¹Kg⁻¹ D. <span style="font-size:
0.5pt; line-height: 107%; font-family: Arial, sans- serif; background-origin: initial; background-clip: initial; '>1.7588 x 10¹¹Kg⁻¹
159	Charge on electron was discovered by:	A. Millikan. B. Crook. C. Neil Bohr. D. Rutherford.
160	Proton was discovered by:	A. Chadwick B. J.J. Rhomson C. Millikan. D. Goldstein.
161	Which one of the following particles has amass 1/1836 time, that of hydrogen?	A. Neutron. B. Proton. C. Electron. D. Positron.
162	Negatively charged particle nature of cathode rays was first demonstrated in 1895 by:	A. Millikan. B. J. Perrin. C. Hittrof D. J.J. Thomson.
163	With the reference of w/m ratio of anode rays, the e/m ratio of cathode rays s:	A. Greater. B. same. C. Smaller. D. Not fixed.
164	Which one of the following statements is true about discovery of neutrons?	 A. These particles were formed by the bombardment of Alpha-particles on Beryllium. B. These particles are formed by the spiting of alpha-particles. C. These particles were discovered by natural radioactivity. D. None of above.
165	Anode is the surface on which probability of finding electron is:	A. 50% B. Less than 10%. C. More than 95%. D. Zero.
166	The experimental evidences for the existence of atomic nucleus comes from:	A. Line spectrum of hydrogen.B. Magnetic bonding of cathode rays.C. Millikan oil drop experiment.D. Scattering of alpha particles by thin metal foil.