

Physics ICS Part 2 Chapter 13 Online MCQ's Test

| Sr | Questions | Answers Choice |
|----|---|---|
| 1 | The conventional current is due to the flow of | A. Atoms and molecules B. Positive charge C. Negative charge D. Bot (b) and (c) |
| 2 | When a pot difference of 4 volt is applied across resistance, 10 J of energy is converted Find charge flows | A. 0.20 C B. 2.5 C C. 5.0 C D. 10.0 C |
| 3 | If a charge Q flows through any cross section of the conductor in time t, the current I is | A. $I=Qt$ B. $I= Q/t$ C. $I= Q*t$ D. $I= Q-t$ |
| 4 | Magnetic effect of current is used | A. To detect a current B. To measure a current C. In electric motor D. All of above |
| 5 | During electrolysis process, density of CuSO_4 solution | A. Remains constant B. Decreased C. Increased D. None of these |
| 6 | For non-ohmic devices, the graph between V and I is | A. Not a straight line B. A straight line C. A curve D. All of above |
| 7 | If there is no fourth band, tolerance is shows as | D. 10% |
| 8 | The resistivity of -----decrease with the increase in temp | A. Gold B. Silver C. Copper D. Silicon |
| 9 | A rheostat can be used as variable resistor as well as a----- | A. Potential divider B. Current divider C. Wheat stone bridge D. Power divider |
| 10 | The condition for the wheatstone bridge to be balanced is given by | D. None of above |
| 11 | e.m.f is the conversion of ----- energy into electrical energy | A. Chemical B. Solar C. Light D. None of these |
| 12 | The product of resistance and conductance is | A. 1 B. Resistivity C. Conductance D. Zero |
| 13 | Terminal potential difference is greater than emf of the cell when | A. Circuit is open B. Circuit is closed C. small battery is charged by bigger battery D. None of these |
| 14 | Unit (S.I) of temperature coefficient of resistivity of a material is | A. K B. K^{-1} C. $^{\circ}\text{C}$ D. K^{-2} |
| 15 | In gas the charge carriers are: | A. Electrons B. Ions C. Both a & b D. None of above |
| 16 | The drift velocity is of order: | A. 10^{-13}m/s B. 10^3m/s C. 10^{-3}m/s D. 10^{-4}m/s |

| | | |
|----|---|--|
| 17 | The free electrons experience force. | <p>A. In direction of -E B. In direction of E C. Both A and B D. All of the above</p> |
| 18 | Heat energy is converted into electrical energy. | <p>A. Solar cells B. thermocouples C. Electric generators D. None of above</p> |
| 19 | The heat produced by passage of current. | <p>A. $H = I^2 R t$ B. $H = IR^2 T$ C. $H = I/Rt$ D. $H = I^2 R t$</p> |
| 20 | The vessel containing the tow electrodes and liquid to known as. | <p>A. Chemical cell B. Volt cell C. Volta cell D. Volta meter</p> |
| 21 | The unit of conductivity is: | <p>A. $\text{Ohm}^{-3} \text{m}^{-1}$ B. Ohm m^{-1} C. Both a and b D. Ohm m^{-1}</p> |
| 22 | Tolerance of "Gold" band. | <p>A. $\pm 10\%$ B. $\pm 5\%$ C. $\pm 15\%$ D. $\pm 20\%$</p> |
| 23 | Thermistor with high - ve temperature coefficient are very accurate for measuring low temperature especially near is. | <p>A. 10 kelvin B. 70 kelvin C. 200 kelvin D. 35 kelvin</p> |
| 24 | The value of maximum output power is? | <p>A. $E/4R$ B. $E^2/4R$ C. $E/4R^2$ D. Non of above</p> |
| 25 | The Kirchhoff 1 st rule is manifestation of: | <p>A. Law of conservation of mass B. Law of Conservation of charge C. Law of conservation of energy D. None of above</p> |
| 26 | The algebraic sum of potential change in a closed circuit is zero. | <p>A. Kirchhoff's 1st rule B. Kirchhoff 2nd rule C. Krichhoff's 3rd rule D. Kirchhoff 4th rule &nbsp;rule</p> |
| 27 | Electric power: | <p>A. $V \times I$ B. $V^2 \times I$ C. V/I D. V/I^2</p> |
| 28 | The color code of "Green" | <p>A. 8 B. 3 C. 5 D. 7</p> |
| 29 | The fraction change in resistance per Kelvin is known as: | <p>A. Temperature coefficient of Resistance B. Coefficient of voltage of change C. Thermal &nbsp;expansion D. All of the above&nbsp;</p> |
| 30 | Semiconductor diodes are called: | <p>A. Ohmic B. non ohmic C. Both a & b D. none of above</p> |
| 31 | The unit of resistance is: | <p>A. Ω B. Ωm C. $\Omega^{-1} \text{m}^{-1}$ D. Ωm^{-1}</p> |
| 32 | Seven resistances are connected as shown in the figures . The equivalent resistance between A and B is: | <p>A. 3Ω B. 4Ω C. 4.5Ω D. 5Ω</p> |
| 33 | Three resistors of resistance R each are combined in various ways, Which of the following cannot be obtained? | <p>A. $3R$ B. $2R/4\Omega$ C. $R/3\Omega$ D. $2R/3\Omega$</p> |

| | | |
|----|---|---|
| 34 | Calculate current in 2 $2R/4\Omega$ resistor. | A. 1 A B. $2R/4\Omega$ C. $R/3\Omega$ D. $2R/3\Omega$ |
| 35 | 106 electrons are moving through a wire per second the current developed is: | A. 1.6×10^{-19} A B. 1 A C. 1.6×10^{-13} A D. 106 A |
| 36 | When a wire is stretched and its radius becomes $r/2$, then its resistance will be | A. $16R$ B. $4R$ C. $2R$ D. 0 |
| 37 | A wire uniform cross-section. A length L and resistance R is cut into two equal parts. The resistivity of each part will be: | A. Doubled B. Halved C. Remain the same D. One fourth |
| 38 | The resistivity of two wires is p_1 and p_2 which are connected in series. If there dimensions are same then the equivalent resistivity of the combination will be: | A. $\frac{p_1 + p_2}{2}$ B. $\frac{1}{\frac{1}{p_1} + \frac{1}{p_2}}$ C. $\frac{p_1 p_2}{p_1 + p_2}$ D. $\frac{p_1 + p_2}{p_1 p_2}$ |
| 39 | The powers of two electric bulbs are 100w and 200w. Which are connected to power supply of 220 V. The ratio of resistance of their filament will be: | A. $1:2$ B. $2:1$ C. $1:3$ D. $4:3$ |
| 40 | Thermocouple is an arrangement of two different metals: | A. To convert heat energy into electrical energy B. To produce more heat C. To convert heat energy into chemical energy D. To convert electrical energy into heat energy |
| 41 | A charged conductor has charge on its. | A. Inner surface B. Outer surface C. Middle surface D. Surrounding space |
| 42 | Ampere second stands for the unit of. | A. Charge B. emf C. energy D. Power |
| 43 | The potential difference between the head and tail of an electrical to | A. 600 Volt B. 700 Volt C. 800 Volt D. 900 Volt |
| 44 | A battery move a charge of 40 C around a circuit at constant rate in 20 Sec. The current will be. | A. 2 A B. 0.5 A C. 80 A D. 8 A |

| | | |
|----|---|--|
| | | D. 800 A |
| 45 | If 1×10^7 electrons passes through a conductor in 1.0 micro second , then the current is. | A. 2 A B. 1.6 A C. 2.6×10^{-6} A D. 1.6×10^{-6} A |
| 46 | Drift velocity of electrons is. | A. 10^{-1} m/s B. 10^{-2} m/s C. 10^{-3} m/s D. 10^{-3} m/s |
| 47 | the current which flows from a point at higher. potential to point at lower potential is called. | A. Electric current B. Conventional current C. Either of these D. None of above |
| 48 | By increasing the temperature of conductor, the flow rate of charges. | A. Increase B. Remains constant C. Decreases D. Changes exponentially |
| 49 | 5 A of current flows through a conductor in 2 minutes, charge in the wire is. | A. 500 C B. 600 C C. 400 C D. 10 C |
| 50 | Heat generated by a 40 W bulb in one hour is. | A. 140 J B. 1440 J C. 14400 J D. 144000 J |
| 51 | The head produced by the passage of current through a resistor is. | A. $H = I^2 R t$ B. $H = I R^2 t$ C. $H = 1/R t$ D. $H = I^2 R / t$ |
| 52 | Magnetic effect of current is used in. | A. Toaster B. Electric iron C. Electric motor D. D.C. Battery |
| 53 | For ohmic device the graph between V and I is. | A. A straight line B. Curve C. Hyperbola D. Parabola |
| 54 | Two resistance of 2 Ohm each are connected in parallel combination equivalent resistance will be. | A. 4 Ohm B. 2 Ohm C. 1 Ohm D. 8 Ohm |
| 55 | One ohm is equal to | A. VC-1 B. CV-1 C. AC-1 D. VA $^{-1}$ |
| 56 | The current flowing through each resistor of equal resistance in parallel combination is. | A. Same B. Different C. Zero D. Infinite |
| 57 | The current through a resistance of 100 Ohm when connecting across a source of 220 V is. | A. 22000 A B. 22 A C. 2.2 A D. 0.45 A |
| 58 | When a wire of length 'l' and resistance R is cut into two equal parts then resistivity of each part. | A. is doubled B. Remains the same C. Is halved D. Is one fourth |
| 59 | Specific resistance of a material depends upon. | A. Length B. Area C. Temperature D. Both A and B |
| 60 | The unit of temperature co efficient of resistivity is. | A. Ohm -m B. K $^{-1}$ C. K D. Ohm |
| 61 | The reciprocal of resistance is called. | A. Capacitance B. Resistance C. Conductance D. Inductance |
| 62 | | A. Resistance B. Resistivity |

| | | |
|----|--|--|
| 62 | mho -m-1 is the unit of. | C. Conductance D. Conductivity |
| 63 | The SI unit of resistivity is. | A. Ohm m-2 B. Ohm m-1 C. Ohm m D. Ohm |
| 64 | Resistivity at a given temperature depends upon. | A. Area of cross section B. Length C. Nature of material of conductor D. Both length and area |
| 65 | A substance having the negative temperature coefficient of resistivity out of the following is. | A. Carbon B. Iron C. Tungsten D. Gold |
| 66 | A certain wire has a resistance R, the resistivity of an other wire of an identical material with the first, except for twice its diameter is. | A. 1/4 R B. 4R C. 2R D. Same as R |
| 67 | Colour codes are used to calculate the. | A. Nature of resistor B. Numerical value of resistance C. Potential difference D. Current |
| 68 | In carbon resistors, then value of Blue colour is. | A. 6 B. 7 C. 8 D. 9 |
| 69 | If the resistance of 500 Ohm have fourth band of silver colour then its upper maximum resistance will be. | A. 600 Ohm B. 550 Ohm C. 450 Ohm D. 400 Ohm |
| 70 | Resistance tolerance for gold colour is. | A. 50% B. 30% C. 20% D. 5% |
| 71 | The thermistors convert changes of temperature into. | A. Light energy B. Electric voltage C. Heat D. Sound |
| 72 | A rheostat can operate as. | A. Amplifier B. Potential divider C. Oscillator D. Transformer |
| 73 | Heat sensitive resistors are called. | A. resistors B. Capacitor C. Thermistors D. Inductors |
| 74 | Resistance tolerance of silver band is. | A. 10% B. 6% C. 7% D. 5% |
| 75 | What is the resistance of carbon resistor which has band brown black brown. | A. 100 Ohm B. 1000 Ohm C. 10 Ohm D. 1.0 Ohm |
| 76 | Which one of the following bulbs has the least resistance. | A. 100 W B. 200 W C. 500 W D. 1000 W |
| 77 | An ideal current source shall have resistance | A. Zero B. Finite but not zero C. Infinite D. Depend upon requirement |
| 78 | Kirchhoff's first rule is the manifestation of the law of conservation of. | A. Mass B. Charge C. Energy D. Momentum |
| 79 | Kirchhoff's voltage rule is a way of stating conservation of. | A. Mass B. Charge C. Energy D. Momentum |
| | | A. Compare emf of two cells |

- A. Compare EMF of two cells
 - B. Detect internal resistance of cell
 - C. Measure P.D.
 - D. All of these
-