

Mathematics ECAT Pre Engineering Online Test

Sr	Questions	Answers Choice
1	The distance of the point (2,3) from y-axis is	A. 2 B. 3 C. 5
2	The distance of the point (2, -3) from x-axis is	A2 B3 C. 2 D. 3
3	The distance of the point (-2,3) from x-axis is	A2 B. 2 C. 3 D. 1
4	Question Image	
5	The distance of the point (2.3) from x-axis is	A. 2 B. 3 C. 5
6	The distance of the point (a,b) from y-axis is	A. a B. b C. a + b
7	The probability to get an odd number in a dice thrown once is	A. 1/2 B. 1/6 C. 1/3 D. 2
8	The distance of the point (a, b) from x-axis is	A. a B. b C. a + b
9	How many arrangements of the letters of the word ADDING can be made	
10	The distance between the points (2, 2) and (3, 3) is	A. 10 C. 5 D. 2
11	The distance between the points (1, 2) and (2, 1) is	A. 3 B. 6
12	How many arrangements of the letter of the word PAKPATTAN can be made	
13	The distance between the points $(0, 0)$ and $(2, 1)$ is	A. 5 C. 0 D. 3
14	The distance between the points $(0, 0)$ and $(1, 2)$ is	A. 5 C. 0 D. 3
15	How many arrangements of the letters of the word PAKISTAN cab be made	
16	The distance between the points (0,0) and (x,y) is	A. x ² + y ² B. x C. y
17	The square of the distance between two points $P(x_1, y_1)$ and $Q(x_2, y_2)$ is	
18	How many arrangements of the letters of the word MATHEMATICS can be made	
19	The number of words that can be formed out of the letters of the word ASSASSINATION is	
20	The distance between two points $P(x_1, y_1)$ and $Q(x_2, y_2)$ is	
21	Question Image	A. 36 B. 360 C. 24 D. 6
22	For all points (x,y) on y-axis	A. x is positive B. x = 0 C. x is negative

		D. y = 0
23	20. 19. 18. 17=	
24	For all points (x,y) on x-axis	A. x is positive B. x is negative C. y = 0 D. y is negative
25	For all points (x,y) in fourth quadrant	A. x > 0, y ⁢ 0 B. x > 0, y > 0 C. x ⁢ 0, y ⁢ 0 D. x ⁢ 0, y > 0
26	Question Image	
27	For all points (x,y) in third quadrant	A. x > 0, y < 0 B. x > 0, y > 0 C. x < 0, y < 0 D. x < 0, y > 0
28	n(n - 1) (n - 2) (n - r + 1) =	
29	For all points (x,y) in second quadrant	A. x > 0, y < 0 B. x > 0, y > 0 C. x < 0, y < 0 D. x < 0, y > 0
30	For all points (x,y) in first quadrant	A. x > 0 , y < 0 B. x > 0 , y > 0 C. x < 0 , y < 0 D. x < 0 , y > 0