ECAT Pre General Science Physics Chapter 2 Vectors and Equilibrium Online Test

Sr	Questions	Answers Choice
1	If x-component of a vector is -3 N and y-component is 3 N, then angle of resultant vector will x-axis is:	A. 45<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° B. 315<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° C. 135<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° D. 225<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">°
2	If two forces of magnitudes 3.5 and 2.5 N act on a body such that the angle between the forces is zero, then magnitude of the resultant will be:	A. 1.0 N B. 6 N C. 3.5 N D. 12 N
3	The magnitude of resultant of three vectors is 3. Its x-component is one, y-component is two, then its z-component is:	A. 0 B. 1 C. 2 D. 3
4	The resultant of two velocities 3 m/sec and 400 cm/sec making an angle 90° with each other is:	A. 20 m/sec B. 5 m/sec C. 3 m.sec D. None of these
5	A force of 5 n is acting Y-axis. Its component along X-axis is:	A. 7 N B. 5 N C. Zero D. 10 N
6	Cosine of an angle is positive in:	A. 2nd quadrant B. 3rd quadrant C. 4th quadrant D. All of these
7	The magnitude of the resultant of two forces may be increased by:	A. Increasing the angle between them B. Decreasing the angle between them C. Drawing a triangle to represent them D. None of these
8	The vector in space has:	A. One component B. Two components C. Three components D. None of these

9	Which of the following is scalar quantity?	A. Electric potential B. Velocity C. Momentum D. Force
10	All trigonometric functions (sine, cosine, tangent etc) are positive in:	A. 1st quadrant B. 2nd quadrant C. 3rd quadrant D. 4th quadrant
11	Two forces each of the magnitude F act perpendicular to each other. The angle made by the resultant force with the horizontal will be:	A. 30<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° B. 45<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° C. 60<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° D. 90<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">°
12	Two forces of 10 N and 8 N are applied simultaneously to a body. the maximum value of their resultant is:	A. 2 N B. - 2 N C. 18 N D. 36 N
13	Two forces each of 10 N act on a body, if the force are inclined at 30° and 60° respectively with x-axis, then x-component of their resultant is:	A. 20 N B. 13.66 N C. 10 N D. 8.66 N
14	When a vector is multiplied by a negative number, its direction:	A. Remains the same B. Changes C. Changes by 180<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° D. None of these
15	An vector of 10 N makes an angle of 45° with x-axis. Angle between its rectangular components with be:	A. 45<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° B. 90<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial, sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">° C. 135<span style="font-size: 10.5pt; line-height: 107%; font-family: Arial,

sans-serif; background-image: initial; background-position: initial; background-size: initial; background-repeat: initial; background-attachment: initial; background-origin: initial; background-clip: initial;">°
D. Zero