

ECAT Pre General Science Online Test

Sr	Questions	Answers Choice
1	A dirty carpet is to be cleaned by heating. This is in according with _____ law of motion.	A. First B. Second C. Third D. None of these
2	A certain force gives an acceleration of 2 m/sec^2 to a body mass 5 kg. The same force would give a 20 kg object an acceleration of:	A. 0.5 m/sec^2 B. 5 m/sec^2 C. 1.5 m/sec^2 D. 9.8 m/sec^2
3	Slope of velocity time graph represents:	A. Acceleration B. Speed C. Torque D. Work
4	In above figures, tell which set of graphs shows that a body is moving with uniform velocity:	A. (i) and (ii) B. (ii) and (iii) C. (iii) and (iv)
5	If the velocity time graph is a straight line parallel to the time-axis, then it means:	A. The body is moving with uniform velocity B. The body is moving with uniform acceleration C. The body is at rest D. None of these
6	The magnitude of the force producing an acceleration of 10 m/sec^2 in a body of mass 500 grams is:	A. 3 N B. 4 N C. 5 N D. 6 N
7	The magnitude of the force producing an acceleration of 10 m/sec^2 in a body of mass 500 grams is:	A. 3 N B. 4 N C. 5 N D. 6 N
8	A body is moving with constant velocity of 10 m/sec in the north-east direction. Then its acceleration will be:	A. 10 m/sec^2 B. 20 m/sec^2 C. 30 m/sec^2 D. Zero
9	A body of mass 5 kg is acted upon by a constant force of 20 n for 7 seconds. The total change in momentum will be:	A. 10 NS B. 100 NS C. 140 NS D. 200 NS
10	When brakes are applied to a fast moving car, the passenger will be thrown:	A. Forward B. Backward C. Downward D. none of these
11	Which one of the following is dimensionless:	A. Acceleration B. Velocity C. Density D. Angle
12	The dimension of linear inertia is:	A. MLT^2 B. ML^0T^{-2} C. ML^0T^0 D. MLT^{-1}
13	A ball is dropped from a height of 4.2 meters. To what height it will rise if there is no loss of KE after rebounding?	A. 4.2 m B. 8.4 C. 12.6 D. None of these
14	A body moving with an acceleration of 5 m/sec^2 started with velocity of 10 m/sec . What will be the distance traversed in 10 seconds?	A. 150 m B. 250 m C. 350 m D. 400 m
15	The short distance between two points direction from its initial point to final point is called:	A. Velocity B. Displacement C. Speed D. None of these

		D. Distance
16	Ethanol (alcohol) is a type of:	A. Electric fuel B. Bio fuel C. Nuclear fuel D. None of these
17	Root out the conventional source of energy:	A. Energy from biomass B. hydroelectric energy C. Geothermal energy D. None of these
18	Biomass includes:	A. Crop residue B. Natural vegetation C. Animal dung D. All of these
19	The consumption source if energy is:	A. Energy from biomass B. Hydroelectric energy C. Geothermal energy D. None of these
20	One KWh is equal to:	A. 3.6×10^{22} J B. 3.6 KJ C. 3.6×10^1 KJ D. 3.6 MJ
21	The velocity given to a body to go out of the influence of earth's gravity is known as:	A. Terminal velocity B. Orbital velocity C. Escape velocity D. None of these
22	When two protons are brought closer potential energy of both of them:	A. Increases B. Decreases C. Remains same D. None of these
23	A body of weight 1 N has a kinetic energy of 1 joule when its speed is:	A. 1.46 m sec^{-1} B. 2.44 m sec^{-1} C. 3.42 m sec^{-1} D. 4.43 m sec^{-1}
24	Tick the conservative force:	A. tension in a string B. Air resistance C. Elastic spring force D. Frictional force
25	Work done along a closed path in a gravitational force is:	A. maximum B. Minimum C. Zero D. Unity
26	If x-component of a vector is -3 N and y-component is 3 N, then angle of resultant vector will x-axis is:	A. 45° B. 315° C. 135° D. 225°
27	If two forces of magnitudes 3.5 and 2.5 N act on a body such that the angle between the forces is zero, then magnitude of the resultant will be:	A. 1.0 N B. 6 N C. 3.5 N D. 12 N

28	The magnitude of resultant of three vectors is 3. Its x-component is one, y-component is two, then its z-component is:	A. 0 B. 1 C. 2 D. 3
29	The resultant of two velocities 3 m/sec and 400 cm/sec making an angle 90° with each other is:	A. 20 m/sec B. 5 m/sec C. 3 m.sec D. None of these
30	A force of 5 n is acting Y-axis. Its component along X-axis is:	A. 7 N B. 5 N C. Zero D. 10 N