

Biology FSC Part 2 Chapter 21 Online MCQ's Test

Sr	Questions	Answers Choice
1	Which tumor is delocalized or has branches other than site of orgin.	A. Benign B. Malignant C. Both D. Apoptosis
2	The tumor which is localized and not transferred to other body parts.	A. Malignant B. Benign C. apoptosis D. Necrosis
3	The laptotene and zygotene lasts for.	A. Few hours B. Few days C. Few weeks D. Few years
4	The chromatin material gets condensed by folding and chromosomes appear as thin thread in mitosis at the beginning of.	A. Inter phase B. Pro phase C. Ana phase D. Meta phase
5	Which one is absent in animal cells.	A. spindle B. Centriole C. Chromatids D. Phragmoplast
6	Each bivalent consists of four.	A. Chromosomes B. Chromatids C. spores D. Chiasmata
7	The most critical phase of mitosis, which ensures equal distribution of chromatids in the daughter cells is.	A. Prophase B. Tele phase C. Meta phase D. Anaphase
8	Contractile ring in cytokinesis is formed by	A. Tubulin B. Actin and Myosin C. Keratin D. Cyclin
9	Chromatin network is visible during	A. Interphase B. Prophase C. Anaphase D. Metaphase
10	Microtubules are composed of protein, tubulin and traces of.	A. DNA B. Glycolipid C. RNA D. Phospholipid
11	The number of sets of microtubules originate from each pair of centriole is.	A. 03 B. 04 C. 05 D. 06
12	Karyokinesis involves division of nucleus and cytokinesis refer to	A. Division of whole cell B. Division of centromere C. Division of cytoplasm D. Division of cell wall
13	During cell division , the nuclear division is called	A. Cytokinesis B. Karyokinesis C. Endomitosis D. Plasmolysis
14	During cell division, the nuclear division is called.	A. Cytokinesis B. Karyokinesis C. Plasmolysis D. Karyotype
15	Tissue culture and cloning seek help through.	A. Mitosis B. Meiosis C. endomitosis D. Karyokinesis

16	The microtubules of mitotic apparatus are composed of protein tubulin nd traces of.	A. DNA B. RNA C. Lipids D. Terpenoids
17	Microtubule are composed of	A. Myosin B. Troponin C. Actin D. Tubulin