

General Math 9th Class English Medium Online Test

Sr	Questions	Answers Choice
1	Question Image 	A. A rational number B. An irrational number C. A complex number D. An integer
2	According to grouped data formula of arithmetic mean:	A. $\frac{\sum f_i x_i}{\sum f_i}$ B. $\frac{\sum f_i}{\sum f_i x_i}$ C. $\frac{\sum f_i x_i}{\sum f_i}$ D. $\frac{\sum f_i}{\sum f_i x_i}$
3	X is called	A. Arithmetic mean B. Mode C. Median D. Group data
4	A running total of class frequency is called	A. histogram B. cumulative frequency C. data D. class interval
5	The table which gives the frequency of each score is called:	A. logarithmic table B. grouped table C. ungrouped table D. frequency table
6	If a group of 5 students get marks from 20% to 30% their frequency will be	A. 10 B. 15 C. 20 D. 5
7	The number of times each value appears in the data is called	A. polygon B. frequency C. histogram D. frequency table
8	The formula used to convert Celsius temperature ($^{\circ}\text{C}$) into Fahrenheit temperature ($^{\circ}\text{F}$) is	A. $^{\circ}\text{C} = \frac{9}{5} (^{\circ}\text{F} - 32)$ B. $^{\circ}\text{C} = \frac{9}{5} (^{\circ}\text{C} + 32)$ C. $^{\circ}\text{C} = \frac{9}{5} (^{\circ}\text{F} + 32)$ D. $^{\circ}\text{C} = \frac{5}{9} (^{\circ}\text{F} - 32)$

The formula used to

A. $^{\circ}\text{F} = \frac{9}{5} (^{\circ}\text{C} + 32)$

9 convert Fahrenheit temperature ($^{\circ}\text{F}$) into Celsius temperature ($^{\circ}\text{C}$) is:

- A. $^{\circ}\text{F} = (9/5 \times ^{\circ}\text{C}) + 32$
B. $^{\circ}\text{F} = (9/5 \times ^{\circ}\text{C}) + 32$
C. $^{\circ}\text{F} = 9/5 (^{\circ}\text{F} - 32)$
D. $^{\circ}\text{F} = 9/5 (^{\circ}\text{F} - 32)$

10 1 miles =?

- A. 1.5 km
B. 1.6 km
C. 6.1 km
D. 1.7 km

11 1. In cartesian plane, vertically line is called

- A. x - axis
B. y - axis
C. point
D. origin

12 In cartesian plane, the horizontal line XOY is called

- A. x - axis
B. y - axis
C. origin
D. co-ordinate axis

13 In cartesian plane point 'O' is called

- A. column
B. row
C. origin
D. axis

14 The co-ordinates of origin are

- A. (1,0)
B. (0,1)
C. (0,0)
D. (1,1)

15 First elements ordered pairs is called

- A. column
B. row
C. point
D. origin